



# BOARD QUESTION PAPER : MARCH 2016

## MATHEMATICS AND STATISTICS

Time: 3 Hours

Total Marks: 80

**Note:**

- i. All questions are compulsory.
- ii. Figures to the right indicate full marks.
- iii. Graph of L.P.P. should be drawn on graph paper only.
- iv. Answer to every new question must be written on a new page.
- v. Answers to both the sections should be written in the same answer book.
- vi. Use of logarithmic table is allowed.

### SECTION – I

**Q.1. (A) Select and write the most appropriate answer from the given alternatives in each of the following sub-questions:** (6) [12]

- i. The negation of  $p \wedge (q \rightarrow r)$  is
 

|   |                                       |
|---|---------------------------------------|
| (A) $p \vee (\sim q \vee r)$                    | (B) $\sim p \wedge (q \rightarrow r)$ |
| (C) $\sim p \wedge (\sim q \rightarrow \sim r)$ | (D) $\sim p \vee (q \wedge \sim r)$   |
- ii. If  $\sin^{-1}(1-x) - 2 \sin^{-1}x = \frac{\pi}{2}$  then  $x$  is
 

|                    |                   |
|--------------------|-------------------|
| (A) $-\frac{1}{2}$ | (B) $1$           |
| (C) $0$            | (D) $\frac{1}{2}$ |
- iii. The joint equation of the pair of lines passing through  $(2, 3)$  and parallel to the coordinate axes is
 

|                            |                            |
|----------------------------|----------------------------|
| (A) $xy - 3x - 2y + 6 = 0$ | (B) $xy + 3x + 2y + 6 = 0$ |
| (C) $xy = 0$               | (D) $xy - 3x - 2y - 6 = 0$ |

**(B) Attempt any THREE of the following:** (6)

- i. Find  $(AB)^{-1}$  if  $A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & -2 & -3 \end{bmatrix}$   $B = \begin{bmatrix} 1 & -1 \\ 1 & 2 \\ 1 & -2 \end{bmatrix}$
- ii. Find the vector equation of the plane passing through a point having position vector  $3\hat{i} - 2\hat{j} + \hat{k}$  and perpendicular to the vector  $4\hat{i} + 3\hat{j} + 2\hat{k}$ .
- iii. If  $\vec{p} = \hat{i} - 2\hat{j} + \hat{k}$  and  $\vec{q} = \hat{i} + 4\hat{j} - 2\hat{k}$  are position vector (P.V.) of points P and Q, find the position vector of the point R which divides segment PQ internally in the ratio 2:1.
- iv. Find  $k$ , if one of the lines given by  $6x^2 + kxy + y^2 = 0$  is  $2x + y = 0$ .
- v. If the lines  $\frac{x-1}{-3} = \frac{y-2}{2k} = \frac{z-3}{2}$  and  $\frac{x-1}{3k} = \frac{y-5}{1} = \frac{z-6}{-5}$  are at right angle then find the value of  $k$ .

**Q.2. (A) Attempt any TWO of the following:** (6)[14]

- i. Examine whether the following logical statement pattern is tautology, contradiction or contingency.  
 $[(p \rightarrow q) \wedge q] \rightarrow p$
- ii. By vector method prove that the medians of a triangle are concurrent.
- iii. Find the shortest distance between the lines  $\vec{r} = (4\hat{i} - \hat{j}) + \lambda(\hat{i} + 2\hat{j} - 3\hat{k})$  and  $\vec{r} = (\hat{i} - \hat{j} + 2\hat{k}) + \mu(\hat{i} + 4\hat{j} - 5\hat{k})$  where  $\lambda$  and  $\mu$  are parameters.

**(B) Attempt any TWO of the following:****(8)**

- i. In
- $\triangle ABC$
- with the usual notations prove that

$$(a - b)^2 \cos^2\left(\frac{C}{2}\right) + (a + b)^2 \sin^2\left(\frac{C}{2}\right) = c^2.$$

- ii. Minimize  $z = 4x + 5y$  subject to  $2x + y \geq 7$ ,  $2x + 3y \leq 15$ ,  $x \leq 3$ ,  $x \geq 0$ ,  $y \geq 0$ . Solve using graphical method.
- iii. The cost of 4 dozen pencils, 3 dozen pens and 2 dozen erasers is ₹ 60. The cost of 2 dozen pencils, 4 dozen pens and 6 dozen erasers is ₹ 90 whereas the cost of 6 dozen pencils, 2 dozen pens and 3 dozen erasers is ₹ 70. Find the cost of each item per dozen by using matrices.

**Q.3. (A) Attempt any TWO of the following:****(6)[14]**

- i. Find the volume of tetrahedron whose coterminus edges are  $7\hat{i} + \hat{k}$ ,  $2\hat{i} + 5\hat{j} - 3\hat{k}$  and  $4\hat{i} + 3\hat{j} + \hat{k}$ .
- ii. Without using truth table show that  
 $\sim(p \vee q) \vee (\sim p \wedge q) \equiv \sim p$
- iii. Show that every homogeneous equation of degree two in  $x$  and  $y$ , i.e.,  $ax^2 + 2hxy + by^2 = 0$  represents a pair of lines passing through origin if  $h^2 - ab \geq 0$ .

**(B) Attempt any TWO of the following:****(8)**

- i. If a line drawn from the point  $A(1, 2, 1)$  is perpendicular to the line joining  $P(1, 4, 6)$  and  $Q(5, 4, 4)$  then find the co-ordinates of the foot of the perpendicular.
- ii. Find the vector equation of the plane passing through the points  $\hat{i} + \hat{j} - 2\hat{k}$ ,  $\hat{i} + 2\hat{j} + \hat{k}$ ,  $2\hat{i} - \hat{j} + \hat{k}$ . Hence find the cartesian equation of the plane.
- iii. Find the general solution of  $\sin x + \sin 3x + \sin 5x = 0$ .

**SECTION – II****Q.4. (A) Select and write the most appropriate answer from the given alternatives in each of the following sub-questions:****(6)[12]**

- i. If the function  
 $f(x) = k + x$ , for  $x < 1$   
 $= 4x + 3$ , for  $x \geq 1$   
 is continuous at  $x = 1$  then  $k =$   
 (A) 7 (B) 8 (C) 6 (D) -6
- ii. The equation of tangent to the curve  $y = x^2 + 4x + 1$  at  $(-1, -2)$  is  
 (A)  $2x - y = 0$  (B)  $2x + y - 5 = 0$   
 (C)  $2x - y - 1 = 0$  (D)  $x + y - 1 = 0$
- iii. Given that  $X \sim B(n = 10, p)$ . If  $E(X) = 8$  then the value of  $p$  is  
 (A) 0.6 (B) 0.7 (C) 0.8 (D) 0.4

**(B) Attempt any THREE of the following:****(6)**

- i. If  $y = x^x$ , find  $\frac{dy}{dx}$ .
- ii. The displacement 's' of a moving particle at time 't' is given by  $s = 5 + 20t - 2t^2$ . Find its acceleration when the velocity is zero.
- iii. Find the area bounded by the curve  $y^2 = 4ax$ , X-axis and the lines  $x = 0$  and  $x = a$ .
- iv. The probability distribution of a discrete random variable X is:

|            |   |    |    |    |    |
|------------|---|----|----|----|----|
| $X = x$    | 1 | 2  | 3  | 4  | 5  |
| $P(X = x)$ | k | 2k | 3k | 4k | 5k |

Find  $P(X \leq 4)$ .



v. Evaluate:  $\int \frac{\sin x}{\sqrt{36 - \cos^2 x}} dx$

**Q.5. (A) Attempt any TWO of the following:**

**(6)[14]**

- i. If  $y = f(u)$  is a differentiable function of  $u$  and  $u = g(x)$  is a differentiable function of  $x$  then prove that  $y = f(g(x))$  is a differentiable function of  $x$  and  $\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx}$ .
- ii. The probability that a person who undergoes kidney operation will recover is 0.5. Find the probability that of the six patients who undergo similar operations.
  - a. None will recover.
  - b. Half of them will recover.

iii. Evaluate:  $\int_0^{\pi} \frac{x}{a^2 \cos^2 x + b^2 \sin^2 x} dx$

**(B) Attempt any TWO of the following:**

**(8)**

- i. Discuss the continuity of the following functions. If the function have a removable discontinuity, redefine the function so as to remove the discontinuity.

$$f(x) = \begin{cases} \frac{4^x - e^x}{6^x - 1}, & \text{for } x \neq 0 \\ \log\left(\frac{2}{3}\right), & \text{for } x = 0 \end{cases} \quad \text{at } x = 0$$

- ii. Prove that:

$$\int \sqrt{a^2 - x^2} dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1}\left(\frac{x}{a}\right) + c$$

- iii. A body is heated at  $110^\circ\text{C}$  and placed in air at  $10^\circ\text{C}$ . After 1 hour its temperature is  $60^\circ\text{C}$ . How much additional time is required for it to cool to  $35^\circ\text{C}$ ?

**Q.6. (A) Attempt any TWO of the following:**

**(6)[14]**

i. Prove that:  $\int_0^{2a} f(x) dx = \int_0^a f(x) dx + \int_0^a f(2a - x) dx$

ii. Evaluate:  $\int \frac{1 + \log x}{x(2 + \log x)(3 + \log x)} dx$

iii. If  $y = \cos^{-1}(2x\sqrt{1-x^2})$ , find  $\frac{dy}{dx}$

**(B) Attempt any TWO of the following:**

**(8)**

- i. Solve the differential equation  $\cos(x + y) dy = dx$   
Hence find the particular solution for  $x = 0$  and  $y = 0$ .
- ii. A wire of length  $l$  is cut into two parts. One part is bent into a circle and other into a square. Show that the sum of areas of the circle and square is the least, if the radius of circle is half the side of the square.
- iii. The following is the p.d.f. (Probability Density Function) of a continuous random variable  $X$ :  

$$f(x) = \begin{cases} \frac{x}{32}, & 0 < x < 8 \\ 0, & \text{otherwise} \end{cases}$$
  - a. Find the expression for c.d.f. (Cumulative Distribution Function) of  $X$ .
  - b. Also find its value at  $x = 0.5$  and  $9$ .